

Reports of *Lipoptena fortisetosa* on dogs and in the environment, and evidence of its widespread establishment in Hungary

Adrienn Gréta Tóth^{a,b}, Attila Bende^c, Sándor Hornok^{d,e}, Zsombor Wagenhoffer^a,
Balázs Szulyovszky^b, Viktória Galla^b, Petra Vöröskői^f, Gergő Keve^{d,e,*}

^a Institute for Animal Breeding, Nutrition and Laboratory Animal Science, University of Veterinary Medicine, Budapest, Hungary

^b Cityvet Veterinary Clinic, Budapest, Hungary

^c Institute of Wildlife Biology and Management, University of Sopron, Sopron, Hungary

^d Department of Parasitology and Zoology, University of Veterinary Medicine, Budapest, Hungary

^e HUN-REN-UVMB Climate Change: New Blood-sucking Parasites and Vector-borne Pathogens Research Group, Budapest, Hungary

^f Private Veterinary Practice, Hungary

ARTICLE INFO

Keywords:

Lipoptena fortisetosa

Deer ked

Louse fly

Cervidae

Hungary

ABSTRACT

Lipoptena fortisetosa is a louse fly of East Asian origin that is considered a potential vector of several pathogenic bacteria and is most commonly associated with deer (Cervidae). The species has been detected in multiple countries in Europe; however, its exact distribution range is unknown. A single individual was detected on a dog at a veterinary clinic in Budapest, Hungary, where the presence of this species had not been confirmed previously. After acquiring information regarding the recent whereabouts of the dog, targeted louse fly collection with insect nets was performed in a forest in Central Transdanubia. Of the 30 flying, unfed specimens of *Lipoptena* spp., 23 were morphologically identified as *L. fortisetosa*. Following this, louse fly collections have been conducted across Hungary during the fall of 2025, including the collection of two additional specimens from a dog. According to these results, *L. fortisetosa* is established in several regions of the country, and this parasite can be effectively transported by dogs. The molecular analyses further confirmed the results, and the distinction between *L. fortisetosa* and *Lipoptena cervi*. The findings draw attention to the potential medical and veterinary significance of the emergence of novel vectors that may have been introduced through animal transport.

1. Introduction

Lipoptena fortisetosa is a louse fly of East-Asian origin, most commonly associated with deer (Cervidae), that is likely to have been introduced to Europe with the sika deer (*Cervus nippon*) (Kurina et al., 2019). Based on the available literature, it can be assumed that *L. fortisetosa* is currently spreading across Europe (Andreani et al., 2020; Gałęcki et al., 2021; Yatsuk et al., 2024). Results from other European countries, including Austria (Jentzsch, 2023), Italy (Andreani et al., 2017, 2020; Kurina et al., 2019), Czech Republic (Zeman, 2013), Poland (Gałęcki et al., 2021), and Estonia (Kurina et al., 2019) suggest the widespread European distribution of the insect. Until October 2025, *Lipoptena cervi* was the only species of the genus *Lipoptena* reported from Hungary (Egri and Rigó, 2014; Papp, 2001). During the review of this manuscript, another author reported the occurrence of four additional specimens from Eastern-Hungary (coordinates: 47.09, 20.77), in a local

journal (Szöllősi-Tóth, 2025). However, these findings were not confirmed by molecular analysis, which would have been beneficial, especially in light of the emergence of recently described species such as *Lipoptena andaluciensis* (González et al., 2024).

Although the original host species of the fly, the sika deer (Yatsuk et al., 2024), is not widespread in the wild in Hungary, one of the largest managed population, comprising approximately 100 individuals, is kept in the Fehérvárcsurgó Game Park. At the turn of the 19th and 20th centuries, several exotic game species, including the sika deer (*Cervus nippon*), were introduced to Hungary. The first individuals were brought to the Fehérvárcsurgó estate in 1910. Later reintroductions of the Dybowski's sika deer (*C. nippon hortulorum*) from the Budapest Zoo and the former Soviet Union contributed to the re-establishment of the population (Farág, 2007; Szabó, 2010).

Given that the sika deer, the original host of the fly, occurs only in a few managed populations in Hungary, the aim of this study was to

* Corresponding author. Department of Parasitology and Zoology, University of Veterinary Medicine, Budapest, Hungary.

E-mail address: keve.gergo@univet.hu (G. Keve).

<https://doi.org/10.1016/j.ijppaw.2025.101161>

Received 17 September 2025; Received in revised form 13 November 2025; Accepted 13 November 2025

Available online 13 November 2025

2213-2244/© 2025 The Author(s). Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

determine whether the presence of a single specimen found in the city centre of Budapest was accidental or originated from an established population within the country.

2. Materials and methods

On 2025.08.26, dog (breed: Hungarian vizsla) visited a veterinary clinic in the city central of Budapest, for the purpose of the removal of a previously diagnosed (on 2025.08.12), solitary, well-circumscribed benign fibroadnexal dysplasia (1.2×1 cm) developed in an area of chronic inflammation on the right forelimb, as well as for tartar removal. During the routine examination before the surgery, the veterinarians noticed a single, engorged, wingless louse fly (Diptera: Hippoboscidae) on its fur. Other than the presence of a single parasite and the previously diagnosed benign tumor, no relevant lesions or abnormalities were observed on the dog's integument, and the animal showed no symptoms suggestive of vector-borne pathogen infection. The parasite was sent to the laboratory of the Department of Parasitology and Zoology, University of Veterinary Medicine Budapest for further examination.

The owner of the dog was surveyed for further information: "Has the dog travelled abroad during the summer? Has the dog visited any forests recently? If yes, where?"

Based on the owners' answers, the dog has not been abroad but visited a forest in Central-Transdanubia (Veszprém County, Hungary), a few days before the surgery. The dog had not been treated with any ectoparasitocides.

The aforementioned forest (coordinates: 46.99, 17.91, situated at an altitude of approximately 311 m above sea level) was visited on 2025.09.06. late afternoon and on 2025.09.07 from morning until afternoon for the purpose of louse fly collection. Flying, unfed specimens were collected using an insect net or specimens from the collectors were removed by hand.

Following these, additional collections have been conducted at various sites across Hungary (the shown coordinates are approximate. No sample collection was conducted in protected areas):

In the Mátra Mountains in Northern Hungary on 2025.09.27 (coordinates: 47.90, 20.01, situated at an altitude of approximately 336 m above sea level).

In the Őrség region in Western Hungary on 2025.10.12 (coordinates: 46.85, 16.51, situated at an altitude of approximately 228 m above sea level).

Two additional louse flies were found by one of the authors after walking a dog in the Bükk Mountains in Northeastern- Hungary, on 2025.10.13 (coordinates: 48.11, 20.61; approximately 319 m above sea level). These specimens, already wingless but not yet engorged, were removed from the dog, which showed no signs of dermatological abnormalities.

The louse flies were identified based on standard morphological keys (Andreani et al., 2019; Maa, 1965; Oboňa et al., 2022; Salvetti et al., 2020) and were also compared to the newly described species *L. andalucensis* (González et al., 2024; Usai et al., 2025). For species identification, the following morphological attributes were used: the size of the specimens (Fig. 2), the number and position of acrostical and postalar bristles (Fig. 3), the mesosternal spines (Fig. 4/A), and the pregenital sclerite of females (Fig. 4/B)).

Four flies (including the *L. fortisetosa* that was initially found on the dog) were chosen for molecular identification: two *L. fortisetosa*, and two *L. cervi*. The surface of louse flies was disinfected by sequential washing for 15 s in detergent, tap water and in distilled water. For the DNA extraction, two legs of each *L. fortisetosa*, and one leg of each *L. cervi* specimens were cut off. DNA was extracted with the QIAamp DNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instruction, including an overnight digestion in tissue lysis buffer and Proteinase-K at 56 °C. Extraction controls (tissue lysis buffer) were also processed with the hippoboscid samples to monitor cross-contamination.

The cytochrome c oxidase subunit I (cox 1) encoding gene was chosen as the target for molecular analysis. The PCR was modified from Folmer et al. (1994) and amplifies an ~710-bp-long fragment of the gene. The primers HCO2198 (5'-TAA ACT TCA GGG TGA CCA AAA AAT CA-3') and LCO1490 (5'-GGT CAACAA ATC ATA AAG ATA TTG G-3') were used in a reaction volume of 25 µl, containing 1 U (stock 5 U/µl) HotStarTaq Plus DNA Polymerase (Qiagen, Hilden, Germany), 2.5 µl 10 × CoralLoad Reaction buffer (including 15 mM MgCl₂), 0.5 µl PCR nucleotide Mix (Qiagen, Hilden, Germany) (stock 10 mM), 0.5 µl of each primer (stock 50 µM), 15.8 µl ddH₂O and 5 µl template DNA. For amplification, an initial denaturation step at 95 °C for 5 min was followed by 40 cycles of denaturation at 94 °C for 40 s, annealing at 48 °C for 1 min and extension at 72 °C for 1 min. Final extension was performed at 72 °C for 10 min.

A non-template reaction mixture served as the negative control in all PCR analyses. Extraction controls and negative controls remained PCR negative in all tests. The PCR products were purified and sequenced by Eurofins Biomi Ltd. (Gödöllő, Hungary). The BioEdit program was used for quality control and trimming of sequences. The analyses of assembled sequences were performed with BLASTN via GenBank (<https://blast.ncbi.nlm.nih.gov>). The sequences obtained in the current study were deposited in the GenBank database and are available under accession numbers PX467064-PX467067.

3. Results

Based on morphological characteristics the single, specimen collected from the dog on 2025.08.26. was identified as *L. fortisetosa* (Fig. 1).

During the louse fly collections on 2025.09.06 and 2025.09.07, 30 specimens of *Lipoptena* spp. were collected. Twenty-three of these were identified as *L. fortisetosa*, while 7 as *L. cervi* (Figs. 2 and 3).

On 2025. 09.27, five specimens were collected, three of which were identified as *L. fortisetosa*, and 2 as *L. cervi*.

On 2025.10.12, 45 louse flies were collected, all of them being *L. cervi*.

Both flies that were removed from the dog on 2025.10.13 were identified as *L. fortisetosa*.



Fig. 1. Wingless, engorged specimen of *Lipoptena fortisetosa*, removed from a dog.



Fig. 2. Unfed specimens of *Lipoptena cervi* (right) and *Lipoptena fortisetosa* (left).

The results of our molecular analyses based on the *cox1* gene confirmed our morphological identifications. The flies we identified as *L. fortisetosa* (PX467064 and PX467065) showed 99.85 % (652/653 bp) identity to a *L. fortisetosa* specimen from Romania (MK405669) and 100 % (643/643 bp) identity to a specimen from Estonia (MN807844). The flies we identified as *L. cervi* (PX467067 and PX467066) showed 100 % (650/650 bp) and 99.85 % identity (649/650 bp) respectively to several *L. cervi* specimens from Scandinavia (KR362270, KR362275).

4. Discussion

The occasional presence of *L. fortisetosa* on dogs has been reported in Europe (Kurina et al., 2019; Mihalca et al., 2019; Sokół and Gałęcki, 2017). In addition to this species, *L. cervi* (Hermosilla et al., 2006), *Hippobosca equina* (Mašlanko et al., 2022) and *Hippobosca longipennis* (Mihalca et al., 2019) are among the most commonly recorded hippoboscids on dogs. In addition, *L. fortisetosa* has been documented parasitizing also various other mammals, including deer, cattle, goats, sheep and humans in Europe (Büttiker, 1994; Kurina et al., 2019; Metelitsa and

Veselkin, 1989; Mihalca et al., 2019; Schledl, 2017). Our report points out, that in contrast to our previous knowledge, dogs are exposed to *L. fortisetosa* in the Hungarian forests, and that the parasite can be passively transported by dogs over considerable distances, contributing to the dissemination of the species. This is supported by the fact that we found this fly on two different dogs. In the initial case, the infested dog had been transported for at least 120 km, and the fly remained on the dog for more than one day, reaching an engorged state. Flies with detached wings were observed on dogs on three occasions (one engorged specimen in Budapest, and two not engorged ones in the Bükk Mountains). This result nuances the previous hypothesis that the species reached Europe through the transportation of deer (Yatsuk et al., 2024), and suggest that pet dogs may play a marginal role in the local dissemination of these ectoparasites. Neither of the dogs exhibited any relevant clinical signs after the removal of the parasites.

Despite the latter facts, it is difficult to determine whether these specimens arrived in Hungary as a result of natural host (deer) migration or through human activity, as, according to new results from Italy, European (including Central European) samples do not differ much from Asian specimens, neither morphologically nor in their molecular characteristics (Andreani et al., 2020). *Lipoptena fortisetosa* was first described in Europe as *L. parvula* in 1967, in what was then Czechoslovakia (Theodor, 1967). This suggests that the previous absence of *L. fortisetosa* from the Hungarian fauna (Egri and Rigó, 2014; Kaufman, 1988; Papp, 2001) could hardly be interpreted as the true absence of the species in the country, since this absence may be explained (for example) by old, but localized and yet undiscovered populations. Obviously, finding an answer to the question: “How and when *L. fortisetosa* arrived in Europe?” is almost impossible; however, the question itself highlights the importance of the parasite monitoring of imported and transported animals.

To the best of our knowledge, this is the first report confirming the presence and establishment of *L. fortisetosa* in Hungary based on both molecular and morphological evidence. Our data indicate that the species now established in several regions in the country. In contrast to previous knowledge, dogs that were walked in forests in central, eastern, and northern Hungary were found to be exposed to this parasite. Furthermore, our findings suggest that *L. fortisetosa* may be unintentionally transported not only by deer but also by dogs.

CRediT authorship contribution statement

Adrienn Gréta Tóth: Writing – review & editing, Writing – original draft, Visualization, Validation, Investigation, Conceptualization. Attila Bende: Writing – review & editing, Writing – original draft,




Fig. 3. The thorax of *Lipoptena cervi* (A) and the thorax of *Lipoptena fortisetosa* (B), with the acrostical and postalar bristles visible in the cases of both species.




Fig. 4. Ventral view of the thorax and abdomen of an unfed female *Lipoptena fortisetosa*. Thorax showing species-specific chaetotaxy of the setae (A). Abdomen displaying the species-specific chaetotaxy of the setae on the first abdominal sternite and the pregenital sclerite (B).

Investigation. **Sándor Hornok:** Writing – review & editing, Investigation. **Zsombor Wagenhoffer:** Writing – review & editing, Investigation. **Balázs Szulyovszky:** Writing – review & editing, Investigation. **Viktória Galla:** Writing – review & editing, Investigation. **Petra Vöröskői:** Writing – review & editing, Investigation. **Gergő Keve:** Writing – review & editing, Writing – original draft, Visualization, Validation, Methodology, Investigation, Formal analysis, Data curation, Conceptualization.

Funding

Financial support was provided by the Office for Supported Research Groups, Hungarian Research Network (HUN-REN), Hungary (Project No. 1500107).

Declaration of competing interest

The authors declare no conflicts of interests.

Acknowledgements

We would like to express our gratitude towards the owner of the infested dog, for her invaluable help.

References

- Andreani, A., Belcari, A., Sacchetti, P., Ponzetta, M.P., 2017. A New Parasites of Ungulates Italy: *L. fortisetosa* (Diptera: Hippoboscidae). *ITA*.
- Andreani, A., Giangaspero, A., Marangi, M., Barlaam, A., Ponzetta, M.P., Roy, L., Belcari, A., Sacchetti, P., 2020. Asia and Europe: so distant So close? The case of *Lipoptena fortisetosa* in Italy. *Kor. J. Parasitol.* 58, 661–668. <https://doi.org/10.3347/kjp.2020.58.6.661>.
- Andreani, A., Sacchetti, P., Belcari, A., 2019. Comparative morphology of the deer ked *Lipoptena fortisetosa* first recorded from Italy. *Med. Vet. Entomol.* 33, 140–153. <https://doi.org/10.1111/mve.12342>.
- Büttiker, W., 1994. Die Lausfliegen der Schweiz (Diptera, Hippoboscidae) mit Bestimmungsschlüssel. Centre suisse de cartographie de la faune, Neuchâtel.
- Egri, B., Rigó, E., 2014. A háság gímszarvasainak *Lipoptena cervi* (Linnaeus, 1758) fertőzőttségéről Irodalmi összefoglaló és saját megfigyelések [on the infestation of Red Deer in the Háság Region by *Lipoptena cervi* (Linnaeus, 1758): Literature review and personal observations. *Magy. Állatorvosok Lapja* 115–122.
- Farágó, S., 2007. Vadászati állattan [game zoology]. Mezőgazda Kiadó 388–393. Budapest.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* 3, 294–299.
- Gałęcki, R., Xuan, X., Bakula, T., Jaroszewski, J., 2021. Molecular characterization of *Lipoptena fortisetosa* from environmental samples collected in North-Eastern Poland. *Animals* 11, 1093. <https://doi.org/10.3390/ani11041093>.
- González, M.A., Ruiz-Arrondo, I., Magallanes, S., Oboña, J., Ruiz-López, M.J., Figuerola, J., 2024. Molecular and morphological analysis revealed a new *Lipoptena* species (Diptera: Hippoboscidae) in southern Spain harbouring *Coxiella burnetii* and bacterial endosymbionts. *Vet. Parasitol.* 332, 110300. <https://doi.org/10.1016/j.vetpar.2024.110300>.
- Hermosilla, C., Pantchev, N., Bachmann, R., Bauer, C., 2006. *Lipoptena cervi* (deer ked) in two naturally infested dogs. *Vet. Rec.* 159, 286–287. <https://doi.org/10.1136/vr.159.9.286>.
- Jentzsch, M., 2023. Die Lausfliegen (Diptera: Hippoboscidae) in der Sammlung des Landesmuseums für Kärnten Klagenfurt. *Entomol. Austriaca* 30, 147–154.
- Kaufman, G., 1988. Magyarország kullancslégy fajai és a fecske-kullancslégy biológiája [The louse fly species of Hungary, and the biology of the swallow-louse fly]. Állatorvostudományi Egyetem, Budapest.
- Kurina, O., Kirik, H., Ounap, H., Öunap, E., 2019. The northernmost record of a blood-sucking ectoparasite, *Lipoptena fortisetosa* Maa (Diptera: Hippoboscidae). *Estonia. Biodivers Data J* 7, e47857. <https://doi.org/10.3897/BDJ.7.e47857>.
- Maa, T.C., 1965. A synopsis of the lipopteninae (Diptera: Hippoboscidae). *J. Med. Entomol.* 2, 233–248. <https://doi.org/10.1093/jmedent/2.3.233>.
- Maślanko, W., Szway, E., Gazda, M., Bartosik, K., 2022. *Hippobosca equina* L. (Hippoboscidae: Hippobosca)—An old enemy as an emerging threat in the palearctic zone. *Int. J. Environ. Res. Publ. Health* 19, 16978. <https://doi.org/10.3390/ijerph192416978>.
- Metelitsa, A.K., Veselkin, G., 1989. Parasitism of the louse fly *Lipoptena fortisetosa* on cattle. *Parazitologia* 23, 276–277.
- Mihalca, A.D., Păstrav, I.R., Sándor, A.D., Deak, G., Gherman, C.M., Sarmaşi, A., Votýpka, J., 2019. First report of the dog louse fly *Hippobosca longipennis* in Romania. *Med. Vet. Entomol.* 33, 530–535. <https://doi.org/10.1111/mve.12395>.
- Oboña, J., Fogašová, K., Fulín, M., Greš, S., Manko, P., Repaský, J., Roháček, J., Sychra, O., Hromada, M., 2022. Updated taxonomic keys for European Hippoboscidae (Diptera), and expansion in central Europe of the bird louse fly *Ornithomya comosa* (Austen, 1930) with the first record from Slovakia. *ZooKeys* 1115, 81–101. <https://doi.org/10.3897/zookeys.1115.80146>.
- Papp, L., 2001. Checklist of the diptera of Hungary. Hungarian Natural History Museum.
- Salveti, M., Bianchi, A., Marangi, M., Barlaam, A., Giacomelli, S., Bertoletti, I., Roy, L., Giangaspero, A., 2020. Deer keds on wild ungulates in northern Italy, with a taxonomic key for the identification of *Lipoptena* spp. of Europe. *Med. Vet. Entomol.* 34, 74–85. <https://doi.org/10.1111/mve.12411>.
- Schledl, W., 2017. Lausfliegen an Cervidae und Bovidae in Tirol (Insecta: diptera, Pupipara, Hippoboscidae). *Linzer boil Beiträge* 49, 885–894.
- Sokół, R., Gałęcki, R., 2017. Prevalence of keds on city dogs in central Poland. *Med. Vet. Entomol.* 31, 114–116. <https://doi.org/10.1111/mve.12209>.
- Szabó, I., 2010. A bakony füttöntgető krakélere [The Whistling Chatterbox of the Bakony]. *Nimród* 98, 19.
- Szöllősi-Tóth, P., 2025. First Record of *Lipoptena fortisetosa* Maa, 1965 from Hungary (Diptera: Hippoboscidae). *Folia Entomologica Hungarica*, pp. 183–186.
- Theodor, O., 1967. *Lipoptena parvula*, n. sp., eine neue Art aus der Tschechoslowakei (Diptera, Hippoboscidae). *Acta Entomol. Musei Natl. Pragae*.
- Usai, F., Dini, F.M., Guarniero, I., Bellinello, E., Stancampiano, L., 2025. The enigmatic case of *Lipoptena* sp. In: The Bosco Della Mesola Nature Reserve (Italy). *Medical Vet Entomology Mve*, 70002. <https://doi.org/10.1111/mve.70002>.
- Yatsuk, A., Triseleva, T., Matyukhin, A., Safonkin, A., Nartshuk, E., 2024. Two cases of introducing *Lipoptena fortisetosa* Maa (Diptera: Hippoboscidae) into Europe through different deer species. *J. Nat. Hist.* 58, 1787–1801. <https://doi.org/10.1080/00222933.2024.2395905>.
- Zeman, P., 2013. Borrelia-infection rates in tick and insect vectors accompanying human risk of acquiring Lyme borreliosis in a highly endemic region in central Europe. *Folia Parasitol.* 45, 319–325.